arXiv:1110.6680v1 [math.NT] 31 Oct 2011

ON A PROBLEM OF BART GODDARD

PING XI

Abstract. We find identities for two series proposed by Bart Goddard in terms of elementary functions.

1. Introduction and the main result

During the 2009 Western Number Theory held at Asilomar, Bart Goddard proposed a problem to ask the expressions of the series

$$A_k(y) := \sum_{n\geqslant 1}(-1)^{n+1}\binom{2n+1}{2k}\frac{y^{2n-1}}{(2n+1)!},$$

$$B_k(y) := \sum_{n\geqslant 1}(-1)^{n+1}\binom{2n+1}{2k+1}\frac{y^{2n-2}}{(2n+1)!}$$

for any fixed natural number k. See [M] [Problem 009:14] for details.

In this short note, we give the desired expressions for $A_k(y)$ and $B_k(y)$. To this end, we consider the following series

$$S_k(y) = \sum_{n\geqslant 1}(-1)^{n+1}\binom{2n+1}{k}\frac{y^{2n+1}}{(2n+1)!}. \tag{1}$$

Clearly, $A_k(y) = y^{-2}S_{2k}(y)$ and $B_k(y) = y^{-3}S_{2k+1}(y)$ for $y \neq 0$. As usual, we regard the series above as the formal power series, and we don't consider their convergence in y.

In general, we can prove that

Theorem 1. *We have*

$$S_0(y) = y - \sin y,$$

$$S_1(y) = y - y\cos y,$$

and for $k \geqslant 2$, *we have*

$$S_k(y) = \frac{y^k}{k!} \times \begin{cases} (-1)^{(k+2)/2}\sin y, & k \text{ even},\\ (-1)^{(k+1)/2}\cos y, & k \text{ odd}. \end{cases}$$

The proof is an easy application of the method of generating functions, which will be outlined in the next section.

The work is supported by N.S.F. (No. 11171265) of P.R. China.

2. PROOF OF THE THEOREM

Now we put

$$S(x,y)=\sum_{k\geqslant 0}S_k(y)x^k. \tag{2}$$

In view of (1) and changing the order of summation, we have

$$S(x,y)=\sum_{n\geqslant 1}(-1)^{n+1}\frac{y^{2n+1}}{(2n+1)!}\sum_{k\geqslant 0}\binom{2n+1}{k}x^k=-\sum_{n\geqslant 1}(-1)^n\frac{(xy+y)^{2n+1}}{(2n+1)!}.$$

Hence we have

$$S(x,y)=xy+y-\sin(xy+y)=xy+y-\sin xy\cos y-\cos xy\sin y.$$

Using the Taylor expansions for $\sin x$ and $\cos x$, we get

$$S(x,y)=\sum_{k\geqslant 0}\beta_k(y)\frac{x^k}{k!}, \tag{3}$$

where $\beta_0(y)=y-\sin y, \beta_1(y)=y-y\cos y$, and

$$\beta_k=\begin{cases}(-1)^{(k+2)/2}y^k\sin y, & k\geqslant 2 \text{ even},\\ (-1)^{(k+1)/2}y^k\cos y, & k\geqslant 2 \text{ odd}.\end{cases}$$

By comparing the coefficients in (2) and (3), we arrive at

$$S_k(y)=\frac{\beta_k(y)}{k!},$$

which completes the proof of the theorem.

REFERENCES

[M] G. Myerson, Western Number Theory Problems — 17 & 19 Dec 2009, http://www.ma.utexas.edu/users/goddardb/WCNT11/problems2009.pdf.

SCHOOL OF MATHEMATICS AND STATISTICS, XI'AN JIAOTONG UNIVERSITY, XI'AN 710049, P. R. CHINA
E-mail address: `xprime@163.com`